

NORTH REGION — CROP MANAGEMENT SUMMARY

The wheat test was conducted at four locations in the North Region (Trenton, Novelty, Columbia, and Martinsburg). The Martinsburg location was managed with higher than normal level of management. Frequent and heavy rains delayed harvest of the Novelty location, but yields and test weights were consistent with other locations.

Characteristics of the wheat test locations

Location	Soil type	Soil test				Precipitation			
		pH	OM %	P (lbs/ac)	K (lbs/ac)	March	April	May	Total
Columbia	Mexico silt loam	6.6	2.0	75	284	2.6	8.5	4.5	15.6
Martinsburg	Calwoods silt loam	5.8	2.4	31	204	3.5	7.9	4.6	16.0
Novelty	Putnam silt loam	5.6	1.5	39	192	3.2	6.2	2.4	11.8
Trenton	Lagonda silty clay loam	5.6	3.0	11	206	3.8	4.8	2.6	11.2

Crop management practices at the wheat test locations

Location	Dates		Fertilizer			Tillage	Herbicides		Fungicide/ Insecticides
	Planting	Harvest	N	P ₂ O ₅	K ₂ O		Pre	Post	
Columbia	Oct. 10	June 13	120	0	0	No Till	Harmony, Roundup	None	Headline
Martinsburg	Oct. 11	June 20	138	100	120	No Till	Harmony, Roundup	Harmony	Prosario
Novelty	Oct. 2	July 8	94	46	60	No Till	None	None	None
Trenton	Oct. 1	June 22	150	0	0	No Till	Harmony	None	Priaxor